

Report 84-43  
Stanford -- KSL

Scientific DataLink

Counterfactuals.  
Matthew L. Ginsberg,  
Dec 1984

card 1 of 1

Stanford Heuristic Programming Project  
Report No. HPP-84-43

December 1984

# Counterfactuals

Matthew L. Ginsberg

COMPUTER SCIENCE DEPARTMENT  
Stanford University  
Stanford, California 94305

# Counterfactuals

## Abstract

Counterfactuals are a form of commonsense non-monotonic inference that has been of long-term interest to philosophers. In this paper, we begin by describing some of the impact counterfactuals can be expected to have in artificial intelligence, and by reviewing briefly some of the philosophical conclusions which have been drawn about them. Philosophers have noted that the content of any particular counterfactual is in part context-dependent; we present a formal description of counterfactuals that allows us to encode this context-dependent information clearly in the choice of a sublanguage of the logical language in which we are working. Having made this choice, we show that our description of counterfactuals is formally identical to the accepted "possible worlds" interpretation due to David Lewis. Finally, we examine the application of our ideas in the domain of automated diagnosis of hardware faults.

## §1. Introduction

A counterfactual is a statement such as, "if  $p$ , then  $q$ ," where  $p$  is expected to be false. Typical examples are, "If the electricity hadn't failed, dinner would have been ready on time," or "If the bedroom door were open, I could get the widget I left in there."

From the point of view of logical semantics, counterfactuals are always true. This is in contrast with our intuitive understanding of their content, which might well accept the statements in the last paragraph while rejecting, for example, "If the power hadn't failed, pigs would fly."

Indeed, the distinction between true and false counterfactuals seems to underly much of our use of knowledge. When planning the solution to a complicated problem, we reduce it to subproblems by realizing that we can prove a counterfactual of the form, "If only thus-and-so were true, I would be able to solve the original problem." The original problem reduces to proving the counterfactual (in some suitable sense) and to arranging for thus-and-so to be true.

Consider the problem of crossing a river if the only boat available has no oars. The counterfactual, "If I had some oars, I'd be able to cross the river," suggests replacing the original problem with that of finding something with which to row. This is a fairly general phenomenon: counterfactuals suggest goal regressions.

Counterfactuals also enable us to understand why plans fail. The example we have already mentioned, "If the electricity hadn't failed, dinner would have been ready on time," is typical. The electricity *did* fail. But in spite of its lack of logical content, the statement does explain why the soup isn't ready.

Dave Smith has pointed out that additional applications can be found in the area of design. Suppose that a machine contains an on-line representation of the design of a complex device. Questions of the form, "If I were to remove the pullup resistor connected to the output of the or gate, would the output of the circuit change?" are counterfactual in nature.

Diagnosis is similar. The counterfactual, "If the device fails in this fashion, the and gate is not functioning," allows us to reason directly from the *intended* description of the design of the device in question, as opposed to reasoning from a description that explicitly allows for the failure of some component, as in [4]. The nature of counterfactual implication is also such as to subsume the single fault assumption—the above counterfactual will be true if (and only if) a fault in the and gate is sufficient, in and of itself, to explain the failure of the device.

Finally, counterfactuals will necessarily play a part in natural language understanding. The extent to which they pervade our communications makes it inevitable that we will eventually need a formal description of them.

My aim in this paper is twofold. Firstly, I would like to describe briefly some of the existing philosophical work that has been done on counterfactuals, although with an eye toward eventual applications in artificial intelligence. Secondly, I will present a formal description of counterfactuals that is precise enough to admit a machine implementation.

Before proceeding, though, I should point out that there are other types of conditionals of potential interest in artificial intelligence. An *indicative conditional* is a statement of the form, "If  $x$  is a bird, then  $x$  can fly." There is some consensus among philosophers [1,5] that the truth value of such a conditional is probabilistic; the reason philosophers draw a distinction between indicative conditionals and other probabilistic statements is

because our confidence in "If  $x$  is a bird, then  $x$  can fly" is a function of the likelihood of a randomly selected bird flying, as opposed to the likelihood of the statement being true for a randomly selected instantiation of  $x$ . The statement will be true for the overwhelming majority of such instantiations, since most of them will not be birds.

I will not be concerned with indicative conditionals here. They have already been considered extensively in the non-monotonic literature [7].

There is also a very loose connection between counterfactuals and causality. In the planning examples we have given, the counterfactual "if  $p$ , then  $q$ " corresponds to " $\neg p$  is a cause for  $\neg q$ ." The electricity failure is the cause of the lateness of the dinner. The lack of oars prevents us from crossing the river.

This connection cannot be pushed too far, however. Suppose that after a MYCIN run, we want to know why the machine asked that a certain clinical test be taken. The response is that, "If the result had been positive, the organism would have been rodlike." This is a useful counterfactual, but the causal connection is from the conclusion to the premise, as opposed to the reverse.

An example where there is no causal connection at all can be found in [6]. Suppose that Olga attends a certain party, but that Boris, who is trying to avoid Olga, does not. If Olga has no similar aversion to Boris, we would have that, "(Even) if Boris had come, Olga would (still) have come." Here, the counterfactual describes the *lack* of a causal connection.

## §2. Properties of counterfactuals

In his excellent book on counterfactuals, Lewis [6] clarifies the distinction between counterfactuals and standard logical implications by listing some of the properties that distinguish them. The results of this section are not new, but may be unfamiliar to an AI audience. A fairly complete treatment of this topic can also be found in [9].

*Contraposition is not a valid rule of inference for counterfactuals.* If we denote the counterfactual, "if  $p$ , then  $q$ ," by  $p > q$ , we cannot conclude  $\neg q > \neg p$  from  $p > q$ . Returning

to our power failure example, it may well be the case that if the power hadn't failed, dinner would have been on time:

*The power didn't fail > dinner was on time.*

It does not follow from this that the electricity *would* have failed if dinner had been late—there may well be an alternative possible cause.

*Counterfactuals are not necessarily transitive.* From  $p > q$  and  $q > r$  we cannot necessarily conclude  $p > r$ . The standard example is due to Stalnaker [10]:

*If J. Edgar Hoover had been born a Russian, then he would have been a Communist, and*

*If he had been a Communist, he would have been a traitor*  
do not together imply

*If Hoover had been born a Russian, he would have been a traitor.*

*Finally, counterfactuals are non-monotonic.* From  $p > r$  we cannot necessarily conclude  $p \wedge q > r$ . In fact, it is possible to have  $p > r$  and  $p \wedge q > \neg r$ : The two statements, "If the electricity hadn't failed, dinner would have been ready on time," and "If the electricity hadn't failed, but I had been elected president, dinner would have been late," are completely consistent.

Glymour and Thomason [5] seem to infer from this last observation that the study of non-monotonic inference generally can be subsumed to some extent under an investigation of counterfactuals, but in light of the breadth of the non-monotonic nature of commonsense reasoning (the frame problem, indicative conditionals, etc.), this seems to me to miss the mark. Counterfactuals in fact seem to be a distinct type of non-monotonic reasoning.

### §3. Possible worlds

Following an idea of Stalnaker's [10], most modern investigations of counterfactuals are based on the notion of possible worlds. Loosely speaking, we analyze a counterfactual  $p > q$  by considering the "possible world" that is as similar to our (real) world as possible,

given that  $p$  is true in it. The counterfactual is true or not depending upon whether or not  $q$  holds in this world.

Lewis [6] has observed that this "most similar possible world" may be unique. He defines a counterfactual to be true if and only if it holds uniformly in the most similar possible worlds in which the premise holds.

This distinction is apparent if we consider the law of the counterfactual excluded middle:

$$(p > q) \vee (p > \neg q).$$

In Stalnaker's view, either  $q$  or  $\neg q$  will hold in the nearest possible world where  $p$  holds, so that the disjunction above will be valid. Lewis points out that this may not be the case by examining the counterfactuals, "If Bizet and Verdi had been compatriots, Bizet would have been Italian," and, "If Bizet and Verdi had been compatriots, Bizet would not have been Italian." Both of these appear to be invalid.

It is possible to understand the discussion of section 2 in terms of possible worlds; we will discuss only the non-monotonic nature of counterfactuals here. The other two properties described earlier are similar.

The basic reason that  $p > r$  and  $p \wedge q > \neg r$  are consistent is that worlds where  $p \wedge q$  hold may be much less similar to our own than worlds where  $p$  alone holds. It is entirely possible that  $r$  holds uniformly in the nearest of the  $p$ -worlds while  $\neg r$  holds uniformly in the (more distant) nearest of the  $p \wedge q$ -worlds. It is also possible that  $r$  holds in some of the nearest of the  $p \wedge q$ -worlds while  $\neg r$  holds in others. In this case we would have only  $p > r$  and  $\neg(p \wedge q > r)$ .

Returning to our power failure example, the nearest possible worlds in which the power remained on are worlds in which dinner was ready on time. In the nearest of the (much) more distant worlds where I was elected president, dinner was late. In still more distant worlds, such as those where the power remained on and I was elected president, but no one bothered to inform me, dinner will once again be prompt.

We can formalize these notions by assuming that, given a fixed world  $F$ , the set of all


possible worlds  $W_\alpha$  can be ordered under a relation  $\geq_F$  where  $U \geq_F V$  if  $U$  is at least as similar to  $F$  as  $V$  is. We can also define  $>_F$  as  $U >_F V$  iff  $\neg(V \geq_F U)$ ; here  $U$  is *more* similar to  $F$  than  $V$  is.

Lewis now defines a counterfactual  $p > q$  to be *true* in some world  $F$  if there exists a world  $U$  such that:

- (1)  $p$  is true in  $U$ , and
- (2)  $p \supset q$  is true in any world  $V$  with  $V \geq_F U$  (i.e., if  $V$  is any world as similar to  $F$  as  $U$  is,  $q$  will hold in  $V$  if  $p$  does).

He also defines  $p > q$  to be (vacuously) *true* if there is no possible world  $U$  in which  $p$  holds.

The following diagram of his may help to explain this. The concentric circles around  $F$  each contain worlds of equal similarity to  $F$ . The diagram on the left corresponds to the case where  $q$  holds uniformly in the most similar of the  $p$  worlds (the shaded region in the diagram), so that  $p > q$  is valid. In the righthand diagram, neither  $q$  nor  $\neg q$  holds uniformly in these worlds, so that neither  $p > q$  nor  $p > \neg q$  is valid.



#### §4. Framework

From an AI perspective, the difficulty with the possible worlds interpretation of counterfactuals is that the notion of "similarity" is too vaguely defined. Our main intention in

this paper is to present a sharper definition of it.

Intuitively, there are (at least) three measures of the similarity, or lack thereof, between different possible worlds. These correspond loosely to the number of propositions whose truth values change, to the extent by which these truth values change, and to the relative importance of the propositions involved.

The last of these is the most difficult to understand in any formal sense, since (as we will see in the next section) it is fundamentally dependent upon context. In fact, we will be able to do little more than to provide a way to encode information concerning the relative importance of the propositions being considered. It is of some interest to note that the scheme we will present can be used to define a notion of similarity that is unchanged from one possible world to another; in other words, we can impose some sort of uniformity on the various relations  $\geq_F$  presented in the last section.

The other two sources of similarity are more syntactic. If the truth value of a proposition changes unnecessarily, in the sense that the possible world without the change is consistent, the possible world which incorporates the change should be deemed "more distant" from our own than the one which doesn't. Similarly, a world where the truth value of some fixed proposition changes from "true" to "true by default" is closer than one where the truth value changes from "true" to "false".

Although this latter is not a feature that will arise in a simple predicate calculus interpretation of counterfactuals, it is important to us because much commonsense reasoning seems to involve these "intermediate" truth values. Since counterfactuals are aspects of commonsense reasoning, they will need to mesh neatly with the features of other commonsense systems.

In order to allow for this, we will work in a logical system that is as loosely defined as possible. While looking for a logical system that potentially includes predicate calculus and some form of default/probabilistic reasoning as special cases, we must also find one capable of supporting precise notions corresponding to logical ideas such as closure and consistency, since these ideas underly that of counterfactual implication. The construction

we are about to present will in fact be a generalization of one due to Ellis [2].

Fix a partially ordered set  $T$  of truth values, and assume that  $T$  admits a least upper bound operation. Denote by  $u$  ("unknown") the least upper bound of all of the elements of  $T$ . Next, let  $L$  be the set of sentences in some language. We define a *truth function* to be a mapping

$$\phi : L \rightarrow T.$$

Intuitively,  $\phi(p) < \phi(q)$  if we are more certain of the truth or falsity of  $p$  than of  $q$ .

Predicate calculus corresponds to  $T = \{u, t, f\}$  ( $t$ =true,  $f$ =false) with  $u > t$  and  $u > f$ ;  $t$  and  $f$  are incomparable. A truth function  $\phi$  assigns to each sentence of  $L$  its truth value, if one is known (and  $u$  otherwise).

We will say that a truth function  $\phi$  is an *extension* of a truth function  $\psi$ , writing  $\phi \leq \psi$ , if  $\phi(p) \leq \psi(p)$  for all  $p \in L$ . We will call an extension  $\phi$  *proper* if the inequality is strict for at least one  $p$  in  $L$ , and will call  $\phi$  a *simple extension* if  $\phi(p) \neq \psi(p)$  for only a single  $p \in L$ .

We will also assume that we have some way of determining whether or not a truth function is consistent. In the predicate calculus case upon which we are modelling our analysis, however, the truth value of a compound sentence is defined recursively in terms of the truth values of its components; furthermore, the definition only applies if the truth values of these components are  $t$  or  $f$  (as opposed to  $u$ ). The (consistent) assignment of the truth values  $t$  or  $f$  to all of the sentences in  $L$  constitutes a *model* for  $L$ .

In general, a model can be characterized by the fact that it is impossible to add to the information corresponding to the truth assignments of the model. Since we have been interpreting  $r < s$  (for truth values  $r$  and  $s$ ) to mean that the information content of the truth value  $r$  is greater than that of  $s$ , it follows that it will be impossible to supplement the information content of the truth assignment  $\phi(p)$  if  $\phi(p)$  is a minimal element of  $T$ .

This leads us to define a truth function  $\phi$  to be a *model* if  $\phi(p)$  is minimal for all  $p \in L$ . If a truth function  $\psi$  is a model that is an extension of the truth function  $\phi$ , we will say that  $\psi$  is a *modular extension* of  $\phi$ . Assuming that consistency is defined as a primitive

for models, we will say that a truth function  $\phi$  is *consistent* iff  $\phi$  has a consistent modular extension.

Here are some predicate calculus examples. In the first two cases,  $\psi$  is a consistent modular extension of  $\phi$ . Since  $\phi$  in the third case has no consistent modular extension, it is inconsistent.

| $x$          | $\phi(x)$ | $\psi(x)$ | $\phi(x)$ | $\psi(x)$     | $\phi(x)$ | $\psi(x)$ |
|--------------|-----------|-----------|-----------|---------------|-----------|-----------|
| $A$          | $t$       | $t$       | $f$       | $f$           | $f$       | $f$       |
| $B$          | $f$       | $f$       | $u$       | $t$ (or $f$ ) | $u$       | $?$       |
| $A \wedge B$ | $t$       | $t$       | $u$       | $f$           | $t$       | $t$       |

**Lemma 1.** Let  $\kappa$  be the set of consistent truth functions. Then if  $\phi \leq \psi$  and  $\phi \in \kappa$ ,  $\psi \in \kappa$ .

**Proof.** Since  $\phi \leq \psi$ , any extension of  $\phi$  is an extension of  $\psi$ .  $\square$

Equivalently, any extension of an inconsistent truth function will also be inconsistent.

We might model default logic with  $T = \{u, t, f, dt, df, pt, pf, *\}$ , with  $dt$ ="true by default",  $df$ ="false by default",  $pt$ ="propensity for truth" and  $pf$ ="propensity for falsity".

Now

$$dt = \text{lub}(t, f, pt) = \text{lub}(t, pt)$$

$$df = \text{lub}(t, f, pf) = \text{lub}(f, pf)$$

$$u = \text{lub}(t, f, pt, pf) = \text{lub}(pt, pf)$$

$$* = \text{lub}(t, f).$$

Probabilistic logic might correspond to  $T = \{[a, b] \subseteq [0, 1]\}$ , with  $[a, b] \leq [c, d]$  if  $[a, b] \subseteq [c, d]$ . The minimal elements of  $T$  in this case are the singletons  $\{x\} \subseteq [0, 1]$ .

For a consistent truth function  $\phi$ , the *closure* of  $\phi$  will be the least upper bound of the consistent modular extensions of  $\phi$ , and will be denoted  $\text{cl}(\phi)$ . If  $\phi = \text{cl}(\phi)$ ,  $\phi$  will be called *closed*. It is not hard to see that this corresponds to logical closure in the predicate calculus case.

In the example below,  $\phi_1$  and  $\phi_2$  are the only consistent modular extensions of  $\phi$ :

|            | $\phi$ | $\phi_1$ | $\phi_2$ | $cl(\phi)$ |
|------------|--------|----------|----------|------------|
| $A$        | $t$    | $t$      | $t$      | $t$        |
| $B$        | $u$    | $t$      | $f$      | $u$        |
| $A \vee B$ | $u$    | $t$      | $t$      | $t$        |

**Lemma 2.**  $cl(\phi) \leq \phi$ .

**Proof.** Let  $\{\phi_i\}$  be the consistent modular extensions of  $\phi$ . Then  $\phi_i \leq \phi$  for all  $i$ , so that  $cl(\phi) = lub\{\phi_i\} \leq \phi$ .  $\square$

Related to the notion of closure is that of reduction: a truth function  $\phi$  will be called *reduced* if all of its simple extensions are consistent. The idea is that a simple extension of  $\phi$  corresponds to the acquisition of more knowledge about some specific proposition; if every such extension is consistent, the original truth function must have been minimal in the sense that it had no extensions which were "necessary" consequences of it.

That the notions of closed and reduced truth functions are not identical is demonstrated by the following example: Returning to the case where  $T$  is the set of closed subintervals of  $[0, 1]$ , suppose that there is a proposition  $p$  such that  $\phi(p) = 0$  is not consistent, but such that  $\phi(p) = x$  is consistent for every  $x \neq 0$ . Now a truth function with  $\phi(p) = [0, 1]$  may well be closed, since  $[0, 1] = lub\{x \in (0, 1]\}$ . It will not be reduced, however, since the simple extension given by  $\phi(p) = 0$  is inconsistent.

In order to formalize the connection between reduction and closure, we need the following definition: The consistency set  $\kappa$  will be called *dense* if, for all collections of truth functions  $\{\phi, \psi_i\}$  which agree uniformly except on a single sentence  $p \in L$ , if  $\psi_i \in \kappa$  for all  $i$ ,  $\psi_i(p)$  is minimal for all  $i$ , and  $\phi(p) \leq lub\{\psi_i(p)\}$ , then  $\phi \in \kappa$ . Intuitively, if  $\kappa$  is dense and the  $\psi_i$  are consistent and agree except at a single proposition  $p$ , then if  $\phi$  is any truth function that agrees with them except at  $p$ , and such that  $\phi(p)$  is in the "span" of the  $\psi_i(p)$ , then  $\phi$  will be consistent as well.

**Lemma 3.** *Predicate calculus is dense.*

**Proof.** If  $\phi(p) \leq lub\{\psi_i(p)\}$  and each  $\psi_i(p)$  is either  $t$  or  $f$ , we must have  $\phi(p) \geq \psi_i(p)$

for some  $i$ . Now apply lemma 1.  $\square$

**Lemma 4.** *Suppose  $\kappa$  is dense. Then any closed truth function  $\phi$  is reduced.*

**Proof.** Let  $\psi$  be a simple extension of  $\phi$  with  $\phi(p) \neq \psi(p)$ . Then there exist consistent modular extensions  $\theta_i$  of  $\phi$  with  $\psi(p) \leq \text{lub}\{\theta_i(p)\}$ . Define truth functions  $\theta'_i$  by

$$\theta'_i(q) = \begin{cases} \phi(q), & \text{if } q \neq p \\ \theta_i(q), & \text{if } q = p. \end{cases}$$

Since  $\theta_i$  is an extension of  $\theta'_i$ , the  $\theta'_i$  are consistent, agree with  $\psi$  except at  $p$ , and  $\psi(p) \leq \text{lub}\{\theta'_i(p)\}$ . Therefore  $\psi$  is consistent.  $\square$

## §5. Similarity

The terminology introduced in the last section allows us to make precise some of the ideas in section 3: possible worlds correspond to consistent models, and sets of possible worlds to consistent truth functions.

The difficulty with this is that we still lack a formal notion of similarity. Part of the problem is context-dependent, as we can see from the pair of counterfactuals

*If Caesar had been in command [in Korea], he would have used the atom bomb*  
and

*If Caesar had been in command, he would have used catapults.*

This example is Quine's [8]. Either counterfactual may well be true (although not both); if the first, Caesar's character is important to our notion of similarity; if the second, it is the tools he had available which are relevant.

It is clearly impossible to select between these two counterfactuals in advance; the best we can do is to present a method for encoding in our semantic machinery the information leading to the choice. In order to do this, let  $L'$  be a subset of  $L$ , and suppose that  $\phi$  is a fixed truth function defined on  $L - L'$ . We can now define a truth function  $\psi$  on  $L'$  to be consistent if and only if the truth function

$$\psi'(p) = \begin{cases} \psi(p), & \text{for } p \in L', \\ \phi(p), & \text{for } p \notin L' \end{cases}$$

is consistent. The effect of this is to fix the truth values outside of  $L'$ , so that any consistent truth function on  $L'$  must be consistent with them.

In the above example, if  $L'$  includes "Caesar was a ruthless military leader," but not, "Caesar's military tools were those of the Roman Empire at its height," we will have to accept Caesar's use of catapults without question, regardless of the weapons available at the time of the engagement, resulting in the validity of the counterfactual which concludes that he would have used catapults. If the situation were reversed, the other counterfactual would be valid. If  $L'$  includes both of the statements describing Caesar, the construction we will present will ambiguously select either of the counterfactuals, while if both of the descriptions are relegated to  $L$ , both counterfactuals will be vacuously true because no truth function  $\phi$  with  $\phi(\text{Caesar in command}) = t$  will have a consistent extension to all of  $L$ .

Given the choice of a (possibly restricted) language  $L'$ , let  $p$  be a sentence in  $L'$ ,  $\phi$  a truth function, and  $s$  a truth value. We will investigate the consequences of the counterfactual premise  $\phi(p) = s$  by defining a new truth function, to be denoted  $\phi|_{\phi(p)=s}$ , and corresponding to  $\phi$  with the truth value at  $p$  replaced by  $s$ . If  $s = t$ , so that we are interested in restricting the value of  $\phi(p)$  to "true", we will denote  $\phi|_{\phi(p)=t}$  simply by  $\phi|_p$ .

We will follow the Caesarian example carefully throughout this section. We have the following propositions in  $L'$ :

$K = \text{Caesar in command in Korea}$

$r = \text{Caesar was ruthless}$

$R = \text{Caesar's tools were those of the Romans}$

$a = \text{the atom bomb was used in Korea}$

$c = \text{catapults were used in Korea.}$

In  $L$  are the rules of inference:

$$K \wedge r \rightarrow a$$

$$K \wedge R \rightarrow c$$

$$\neg(a \wedge c).$$

The last of these is equivalent to  $\neg(K \wedge r \wedge R)$ . Our initial truth assignment is given by:

|     |        |
|-----|--------|
|     | $\phi$ |
| $K$ | $f$    |
| $r$ | $t$    |
| $R$ | $t$    |
| $a$ | $f$    |
| $c$ | $f$    |

Of interest to us is the truth function  $\phi|_K$ ; what if Caesar had been in command in Korea?

The general difficulty is that simply changing  $\phi(p)$  to  $s$  may well produce an inconsistent truth function. We begin therefore by replacing  $\phi(p)$  not with  $s$ , but with the least upper bound of  $s$  and  $\phi(p)$ . This truth function  $\phi'$  must be consistent (since  $\phi$  is an extension of it), but need not be closed.

|     | $\phi$ | $\phi'$                  | $\text{cl}(\phi')$ |
|-----|--------|--------------------------|--------------------|
| $K$ | $f$    | $u = \text{lub}\{t, f\}$ | $f$                |
| $r$ | $t$    | $t$                      | $t$                |
| $R$ | $t$    | $t$                      | $t$                |
| $a$ | $f$    | $f$                      | $f$                |
| $c$ | $f$    | $f$                      | $f$                |

Assume next that there is at least one reduced truth function  $\psi$  of which  $\phi'$  is an extension. (By lemma 3, we will generally be able to take for  $\psi$  the least upper bound of all consistent models for  $L$ .) The set of all such  $\psi$ 's is partially ordered under extension; let  $\phi''$  be a minimal element of it.

|     | $\phi$ | $\phi'$ | $\psi$ (typical) | $\phi''_1$ | $\phi''_2$ |
|-----|--------|---------|------------------|------------|------------|
| $K$ | $f$    | $u$     | $u$              | $u$        | $u$        |
| $r$ | $t$    | $t$     | $t$              | $t$        | $u$        |
| $R$ | $t$    | $t$     | $u$              | $u$        | $t$        |
| $a$ | $f$    | $f$     | $u$              | $u$        | $f$        |
| $c$ | $f$    | $f$     | $u$              | $f$        | $u$        |

The result of replacing  $\phi''(p)$  with  $s$  is necessarily consistent, since  $\phi''$  is reduced and  $\phi''(p) \geq s$  initially. Let  $\theta$  be the truth function obtained by making this replacement, and take  $\phi|_{\phi(p)=s}$  to be  $\text{cl}(\theta)$ . We also define  $\phi(p > q) = \phi|_p(q)$ .

|         | $\phi''_1$ | $\theta_1$ | $\phi _K$ | $\phi''_2$ | $\theta_2$ | $\phi _K$ |
|---------|------------|------------|-----------|------------|------------|-----------|
| $K$     | $u$        | $t$        | $t$       | $u$        | $t$        | $t$       |
| $r$     | $t$        | $t$        | $t$       | $u$        | $u$        | $f$       |
| $R$     | $u$        | $u$        | $f$       | $t$        | $t$        | $t$       |
| $a$     | $u$        | $u$        | $t$       | $f$        | $f$        | $f$       |
| $c$     | $f$        | $f$        | $f$       | $u$        | $u$        | $f$       |
| $K > a$ |            |            | $t$       |            |            | $f$       |
| $K > c$ |            |            | $f$       |            |            | $t$       |

The construction we have given works through the construction of the "intermediate" truth function  $\phi''$ , which is in some sense a set of the most similar possible worlds in which we might have  $\phi(p)$  being either its original value or the new value  $s$ . The reason we have done this is because it seems to be technically easier to define similarity for a world of which ours is an extension ( $\phi'$  in the above construction) than for one which is not directly comparable with it (as  $\phi$  with  $\phi(p)$  replaced by  $s$  would be).

Having constructed similar worlds where  $\phi(p) = s$  might hold, it is straightforward to investigate the consequences if we assume that it *does* hold. This is the content of our taking  $\phi|_{\phi(p)=s}$  to be the closure of  $\phi''$  with  $\phi''(p)$  replaced by  $s$ .

The choice of truth function  $\phi''$  corresponds to choosing a *subset* of the possible worlds in which  $\phi(p) = s$ . In our military example, assuming Caesar to be in command in Korea required our abandoning one of these descriptions, and the choice of which to discard corresponded to our choice of  $\phi''$ .

An alternative would be to assume that the worlds where we abandon one description are just as similar to our own as those in which we abandon another. Lewis might well make this choice; it corresponds to defining

$$\phi(p > q) = \text{lub}\{\phi|_r(q)\},$$

where the least upper bound is taken over all possible choices of  $\phi''$ .

## §6. Formal description

Previous analyses of counterfactuals have generally developed a set of axioms which counterfactual implication must satisfy. We will conclude by showing that our definition

matches Lewis' in the predicate calculus case and with the definition of  $\phi(p > q)$  at the end of the last section.

Lewis introduces two counterfactual operators. He writes  $p \Box \rightarrow q$  for, "If it were the case that  $p$ , then it would be the case that  $q$ ", and  $p \Diamond \rightarrow q$  for, "If it were the case that  $p$ , then it might be the case that  $q$ ." Both of these are handled by our formalism. We write

$$\begin{aligned} p \Box \rightarrow q & \text{ iff } \phi(p > q) = t \\ p \Diamond \rightarrow q & \text{ iff } \phi(p > q) \geq t. \end{aligned}$$

**Theorem 5.**  $p \Box \rightarrow q$  is equivalent to  $\neg(p \Diamond \rightarrow \neg q)$ .

**Proof.** Suppose  $p \Box \rightarrow q$ . Then if  $\psi$  is a consistent modular extension of any  $\theta$  in our counterfactual construction with  $\psi(p) = t$ ,  $\psi(q) = t$ . Thus  $\psi(\neg q) = f$ , so  $\phi(p > \neg q) = f$  and  $\neg(p \Diamond \rightarrow \neg q)$ . The reverse implication is identical.  $\square$

Lewis in fact takes  $\Box \rightarrow$  as primitive and uses theorem 5 to define  $\Diamond \rightarrow$ .

Our main result depends on one due to Gärdenfors [3]:

**Theorem 6 (Gärdenfors).** A counterfactual operator  $\Box \rightarrow$  is formally identically to Lewis' iff it satisfies:

$$(p \Box \rightarrow q) \wedge (p \Box \rightarrow r) \rightarrow (p \Box \rightarrow q \wedge r) \tag{1}$$

$$p \Box \rightarrow \top \tag{2}$$

$$(p \Box \rightarrow q) \wedge \models (q \rightarrow r) \rightarrow (p \Box \rightarrow r) \tag{3}$$

$$p \Box \rightarrow p \tag{4}$$

$$\models (p \leftrightarrow q) \rightarrow ((p \Box \rightarrow r) \rightarrow (q \Box \rightarrow r)) \tag{5}$$

$$(p \wedge q) \rightarrow (p \Box \rightarrow q) \tag{6}$$

$$(p \Box \rightarrow q) \rightarrow (p \rightarrow q) \tag{7}$$

$$(p \Box \rightarrow r) \wedge (q \Box \rightarrow r) \rightarrow (p \vee q \Box \rightarrow r) \tag{8}$$

$$(p \Box \rightarrow q) \wedge \neg(p \Box \rightarrow \neg r) \rightarrow (p \wedge r \Box \rightarrow q). \quad \square \tag{9}$$

**Theorem 7.** *The counterfactual construction of section 5 is formally identical to Lewis' possible worlds construction.*

**Proof.** We need to verify (1) through (9) above. As in the proof of theorem 5, let  $\psi$  be a consistent modular extension of some  $\theta$  in the counterfactual construction. Then:

For (1), we must show that if  $\psi(p) = t$  implies  $\psi(q) = t$  and  $\psi(r) = t$ , it implies  $\psi(q \wedge r) = t$ . This is immediate.

For (2), we must show that  $\psi(\top) = t$ . This is also immediate.

For (3), we must show that if  $\psi(p) = t$  implies  $\psi(q) = t$ , and if  $\chi(q \rightarrow r) = t$  for any consistent model  $\chi$ , then  $\psi(p) = t$  implies  $\psi(r) = t$ .

(4) requires that if  $\psi(p) = t$ , then  $\psi(p) = t$ . For (5), if  $\psi(p) = t$  is equivalent to  $\psi(q) = t$ , then clearly  $\psi(p) = t$  will imply  $\psi(r) = t$  iff  $\psi(q) = t$  does.

In (6), suppose that  $(p \wedge q)$  holds for our original truth function  $\phi$ . Then we will clearly have  $\phi|_p = \phi$ , so that  $p \Box \rightarrow q$  because  $\phi(q) = t$ .

For (7), if  $\phi(p \rightarrow q) \neq t$ , there is a consistent modular extension  $\chi$  of  $\phi$  with  $\chi(p) = t$  and  $\chi(q) = f$ . This  $\chi$  will also be a consistent modular extension of a  $\theta$  in the counterfactual construction, so that  $\phi(p > q) \neq t$ .

The only non-trivial case in (8) is that where  $\phi(p) = \phi(q) = f$ . Now the premise is equivalent to the assumption that in any consistent modular extension  $\chi$  of  $\theta$  where  $\chi(p) = t$  or  $\chi(q) = t$ ,  $\chi(r) = t$ . Thus if  $\chi(p \vee q) = t$ ,  $\chi(r) = t$ .


In (9), suppose first that  $\phi(p) = t$ . Then we know that  $\phi(q) = t$ , and  $\phi(r) \geq t$ . It follows that there is a consistent modular extension  $\chi$  of  $\phi$  with  $\chi(p \wedge r) = t$ , so that  $\phi|_{p \wedge r}$  is an extension of  $\phi$ , and  $\phi|_{p \wedge r}(q) \leq \phi(q) = t$ .  $\phi(p) = u$  is similar, and if  $\phi(p) = f$ , we have that there is a consistent modular extension  $\chi$  of some  $\theta$  with  $\chi(p \wedge r) = t$ , and that for any consistent modular extension of a  $\theta$  with  $\chi(p) = t$ ,  $\chi(q) = t$ . The conclusion follows.  $\square$

## §7. Example: diagnosis

### 7.1 Setting

Genesereth has proposed [4] that it is possible for machines to be used in automated

diagnosis, provided that the machines are given both a design for the device in question, and the ability to manipulate the device by varying its inputs and observing the results. He investigates the diagnosis of a full adder in considerable detail:



A full adder is essentially a one bit adder with carry in and carry out, and it is usually used as one of  $n$  elements in an  $n$  bit adder. A graphical representation of its design is given [above]. It has three inputs and two outputs and consists of two "xor" gates (X1 and X2), two "and" gates (A1 and A2), and an "or" gate (O1) ... In normal operation, the first output (the "sum" line) is "on" if and only if an odd number of inputs is "on"; the second output (the "carry" line) is "on" if and only if at least two inputs are "on". [4]

Genesereth proceeds to describe the design of the full adder in a variant of prefix predicate calculus. We continue to quote him:

```
SD1: (XORG X1)
SD2: (XORG X2)
SD3: (ANDG A1)
SD4: (ANDG A2)
SD5: (ORG O1)

SD6: (CONN (IN 1 F1) (IN 1 X1))
SD7: (CONN (IN 1 F1) (IN 1 A1))
SD8: (CONN (IN 2 F1) (IN 2 X1))
SD9: (CONN (IN 2 F1) (IN 2 A1))
SD10: (CONN (IN 3 F1) (IN 2 X2))
SD11: (CONN (IN 3 F1) (IN 1 A2))
SD12: (CONN (OUT 1 X1) (IN 1 X2))
SD13: (CONN (OUT 1 X1) (IN 2 A2))
SD14: (CONN (OUT 1 A1) (IN 2 O1))
```

SD15: (CONN (OUT 1 A2) (IN 1 O1))  
SD16: (CONN (OUT 1 X2) (OUT 1 F1))  
SD17: (CONN (OUT 1 O1) (OUT 2 F1))

These axioms describe the *structural description* of the full adder. SD1, for example, states that X1 is an exclusive-or gate. SD13 states that the first output of X1 is connected to the second input of A2.

Genesereth also states in a similar form results describing the behavior of the various sorts of gates, and describing in these terms what it means for two points in the circuit to be connected. Using these additional axioms, it is possible to prove that if, for example, the first input to the full adder is on while the other two are off, the first output should be on while the second should be off.

## 7.2 Diagnosis using predicate calculus

The situation of interest, of course, is that in which the outputs of the adder are not as predicted by the design. Genesereth assumes that we have:

AC1: (VAL (IN 1 F1) 1 ON)  
AC2: (VAL (IN 2 F1) 1 OFF)  
AC3: (VAL (IN 3 F1) 1 OFF)  
OB1: (VAL (OUT 1 F1) 1 OFF)  
OB2: (VAL (OUT 2 F1) 1 OFF)

In other words, the values of the three inputs to the adder at time  $t = 1$  are as described at the end of the last subsection, but both of the outputs are off at that time.

The observed behavior characterized by AC1-AC3 and OB1-OB2 is inconsistent with the design given by SD1-SD17. Diagnosis is a matter of resolving this inconsistency.

To do so using predicate calculus, Genesereth assumes (correctly!) that the device in question does not satisfy the design description given earlier, but instead satisfies some weaker "device assumptions". In the example we are considering, he assumes that:

- (1) The connections are all as described in the design, and
- (2) At most one of the gates is broken (the *single fault assumption*).

These device assumptions can be written as:

DA1: (CONN (IN 1 F1) (IN 1 X1))

DA2: (CONN (IN 1 F1) (IN 1 A1))  
 DA3: (CONN (IN 2 F1) (IN 2 X1))  
 DA4: (CONN (IN 2 F1) (IN 2 A1))  
 DA5: (CONN (IN 3 F1) (IN 2 X2))  
 DA6: (CONN (IN 3 F1) (IN 1 A2))  
 DA7: (CONN (OUT 1 X1) (IN 1 X2))  
 DA8: (CONN (OUT 1 X1) (IN 2 A2))  
 DA9: (CONN (OUT 1 A1) (IN 2 O1))  
 DA10: (CONN (OUT 1 A2) (IN 1 O1))  
 DA11: (CONN (OUT 1 X2) (OUT 1 F1))  
 DA12: (CONN (OUT 1 O1) (OUT 2 F1))  
 DA13: (IF (NOT (XORG X1)) (AND (XORG X2) (ANDG A1) (ANDG A2) (ORG O1)))  
 DA14: (IF (NOT (XORG X2)) (AND (XORG X1) (ANDG A1) (ANDG A2) (ORG O1)))  
 DA15: (IF (NOT (ANDG A1)) (AND (XORG X1) (XORG X2) (ANDG A2) (ORG O1)))  
 DA16: (IF (NOT (ANDG A2)) (AND (XORG X1) (XORG X2) (ANDG A1) (ORG O1)))  
 DA17: (IF (NOT (ORG O1)) (AND (XORG X1) (XORG X2) (ANDG A1) (ANDG A2)))

The reappearance of the connection information in DA1-DA12 amounts to the assumption that the connections are intact; the axioms DA13-DA17 encode the single fault assumption.

The axioms above are consistent with the observed behavior of the adder, and lead to the conclusions that

$$(OR (NOT (XORG X1)) (NOT (XORG X2))) \quad (10)$$

and

$$(AND (ANDG A1) (ANDG A2) (ORG O1)). \quad (11)$$

In other words, one of the exclusive-or gates is broken, and the remaining components are functioning satisfactorily. The information about the functionality of the and and or gates is useful because it enables us to generate a test to determine which of the two exclusive-or gates is in fact faulty.

The difficulty with this approach is that it requires us to generate the device assumptions DA1-DA17. It is possible that the fault(s) in the device are such that these assumptions are invalid, in which case the system will be unable to diagnose the device without a replacement set of device assumptions.

### 7.3 Diagnosis using counterfactuals

The device assumptions in the last subsection were introduced to encode our belief that the wiring in the adder was correct, and to enable us to take advantage of the simplifying assumption that only one of the remaining components was damaged. Both of these conditions can be recast naturally in the framework of counterfactuals.

To perform the diagnosis using the methods we have presented, we will assume the structural description DS1-DS17 and the inputs achieved by AC1-AC3, and examine the counterfactual consequences of the observed outputs OB1-OB2. There are three possible explanations for the fault:

- (1) One or more of the components is faulty.
- (2) The wiring is faulty.
- (3) The inputs were not as expected.

We eliminate all but the first from our counterfactual reasoning by including only the component assumptions SD1-SD5 in our restricted language  $L'$ . The wiring and input assumptions SD6-SD17 and AC1-AC3 are therefore assumed to be independent of the counterfactual assumptions corresponding to the observed misbehavior of the device.

The conclusion (10) that one of the two exclusive-or gates must be faulty is in fact a *logical* consequence of the behavior of the device:

$$OB1 \wedge OB2 \rightarrow (OR (NOT (XORG X1)) (NOT (XORG X2))).$$

Meanwhile, because the remaining components cannot be contributing to the observed fault, their continued performance is *counterfactually* implied by the observed behavior:

$$OB1 \wedge OB2 > (AND (ANDG A1) (ANDG A2) (ORG O1)).$$

This reappearance of (11) is especially useful. Because of the non-monotonic nature of counterfactual reasoning, it is of course possible that additional observations appended to the lefthand side of the above equation will invalidate its conclusion; this will happen whenever the single fault assumption is violated. In this case, however, rather than

generating a contradiction, the counterfactual analysis will automatically produce a new diagnosis which once again involves failure for a minimal set of components.

It is possible, however, for a counterfactual analysis to suggest a violation of the single fault assumption when one is not required. If the observed behavior can be explained either by the failure of a single component, or by the failure of a pair of different components, both will be proposed. There is nothing counterintuitive about this, however—it is quite normal to assume that a group of normally undependable components has failed before questioning a single part of proven reliability. In any event, we can if necessary retain the single fault assumption by using it to select among the possible  $\phi$ 's in the counterfactual construction itself.

## §8. Conclusion

Our aim in this paper has been to present a formal description of counterfactuals, describing them in terms of existing logical operators instead of following the usual practice of developing a "counterfactual calculus" to describe their behavior.

We have also sought to work in a setting that will not eventually commit us to any particular reasoning paradigm; since counterfactuals are a form of commonsense reasoning, and there are currently a variety of competing paradigms for commonsense reasoning generally, this seemed prudent.

The construction we have presented seems to meet these objectives. It has indeed described counterfactuals in terms of existing logical primitives, and, in the setting of predicate calculus, reduces to the "possible worlds" interpretation of counterfactuals that is accepted by philosophers.

Our construction also distinguishes clearly between the context dependent and context independent features of counterfactual implication. It provides us with a precise method for selecting those aspects of our world which are to be considered inviolable even under a counterfactual assumption; having made such a choice, we proceed to generate possible worlds which respect it.

The biggest difficulty with the approach we have described is the rather heavy-handed nature of the choice described in the last paragraph. Although it is possible to clearly recognize ambiguities remaining in the analysis of any particular counterfactual (they correspond to the choice of  $\phi''$  in our construction), we have no method for choosing consistently between them. In any specific implementation, it will of course be possible to select an  $\phi''$  when one is needed, but we have not considered the nature of the formalism that should govern this choice.

### Acknowledgement

This research has been supported by the Office of Naval Research under grant number N00014-81-K-0004. A preliminary version of this research was presented to MUGS at Stanford, and the author would like to thank the participants in that seminar for their useful comments on this work. Mike Genesereth and Dave Smith were especially helpful.

### References

- [1] Adams, E., The logic of conditionals, *Inquiry* 8 (1965), 166-197.
- [2] Ellis, B., Epistemic foundations of logic, *J. Phil. Logic* 4 (1975), 133-153.
- [3] Gärdenfors, P., Conditionals and changes of belief, *Acta Philosophica Fennica* 30 (1978), 381-404.
- [4] Genesereth, M.R., The use of design descriptions in automated diagnosis, Stanford Heuristic Programming Project memo HPP-81-20 (1984)
- [5] Glymour, C., and Thomason, R.H., Default reasoning and the logic of theory perturbation, *Non-monotonic Reasoning Workshop*, American Association for Artificial Intelligence (1984) 93-102.
- [6] Lewis, D., *Counterfactuals*, Harvard University Press, Cambridge (1973).
- [7] Perlis, D., Bibliography of literature on non-monotonic reasoning, *Non-monotonic Reasoning Workshop*, American Association for Artificial Intelligence (1984) 396-401.
- [8] Quine, W.V., *Methods of Logic*, Holt, Rinehart, and Winston, New York (1950).
- [9] Sobel, J.H., Utilitarianisms: simple and general, *Inquiry* 13 (1970) 394-449.
- [10] Stalnaker, R., A theory of conditionals, in *Studies in Logical Theory*, Rescher, N. (ed.), Oxford University Press, Oxford (1968).

Copyright © 1985 by KSL and  
Comtex Scientific Corporation

7

FILMED FROM BEST AVAILABLE COPY